

CASE REPORT

PRESSURE SORE

GENERAL INFORMATION

87-year old female patient with a post-surgical pressure sore (grade III), bottom

PREVIOUS TREATMENT

Healing by secondary intention with moist wound care during 30 days without improvement of the wound situation. No systematic pressure relief. Start treatment with ① 30 days after incidence and initial surgical debridement.



DAY 1 Start treatment with ①

TREATMENT WITH ①

WOUND CLEANSING

Saline solution

SECONDARY DRESSING

Non-woven gauze (also used as wound filler)

OTHER

Pressure relief: Not systematically, mobilization of patient.



DAY 21



DAY 49



DAY 71

RESULT

TIME TO WOUND CLOSURE

71 days

OBSERVATIONS

- Clean and vital wound after 2 weeks
- Fast granulation despite lack of systematic pressure relief

CONCLUSION MEDICAL DOCTOR

Dr. F Castrignano, MD
Italy

“Due to a lack of financial resources a surgery could not be performed. „1“ offered a very effective alternative.”

“1” enabled a very simple dressing change that could be performed by the home care organization at home.”

«In addition the patient was spared an additional surgical intervention. This is especially valuable for patients with a high operation risk.»

CONTACTS

PRODUCER

Phytoceuticals AG
CH-8134 Adliswil

FURTHER INFORMATION

www.wound.ch
Tel. +41 58 800 58 58

