

CASE REPORT

TRAUMATIC FINGERTIP AMPUTATION

GENERAL INFORMATION

40-year old male patient with a 4mm deep and 12mm long wound following a fingertip amputation caused by a machine

PREVIOUS TREATMENT

Healing by secondary intention and start treatment with ① on the day of the injury. Wound was initially disinfected.

TREATMENT WITH ①

WOUND CLEANSING

Saline solution

SECONDARY DRESSING

- Non-woven gauze
- Fingerbandage
- Splint

OTHER

Pain medication for 1 week following the injury.

RESULT

TIME TO WOUND CLOSURE

40 days

OBSERVATIONS

The patient or his wife were able to perform the dressing change.



DAY 1 Start treatment with ①



DAY 7



DAY 49



DAY 40

CONCLUSION MEDICAL DOCTOR

Dr. M. Fischer, MD
Switzerland

“The application of “1” resulted in a reduced formation of scab which had a positive effect on the induction of granulation tissue formation.”

“The newly formed granulation tissue cracked open less because “1” maintained a balanced moist wound environment. This had a positive effect on the functionality of the finger.”

“The patient was especially impressed by the fact that he was able to perform the dressing change himself because of the simple application “1” enables.”

CONTACTS

PRODUCER

Phytoceuticals AG
CH-8134 Adliswil

FURTHER INFORMATION

www.wound.ch
Tel. +41 58 800 58 58

