

CASE REPORT


DOG-TRAUMATIC WOUND

GENERAL INFORMATION

Bulldogge, male, 5 years

Dog bite wound with exposed tendon

PREVIOUS TREATMENT

Antibiotic and antiseptic treatment during 1 week without improvement of wound condition. Treatment start with  on day 7 after injury.

TREATMENT WITH

WOUND CLEANSING

Saline solution

SECONDARY DRESSING

None. Dog was wearing collar.

OTHER

Stop antibiotic and antiseptic treatment on day 7 after injury.

RESULT

TIME TO WOUND CLOSURE

22 weeks
(retraumatisation in week 8)

TIME TO HAIR GROWTH

21 weeks

OBSERVATIONS

Quick reduction of itching.



DAY 0 Start treatment with 



DAY 41



DAY 104



DAY 145

CONCLUSION VETERINARY

Dr. Fiorella Carnevali
Italy

"The dressing change was difficult due to the aggression of the dog. By using one, the time for dressing change was significantly shorter, which was very helpful."

"I observed a strong induction of granulation tissue which is impressive given the localisation of the wound."

"As soon as I treated the wound with one the pain and itching seemed to disappear quickly which had a very positive impact on the dog's behaviour."

CONTACTS

PRODUCER

Phytoceuticals AG
CH-8008 Zurich

ADDITIONAL INFO

www.onewound.com
Tel. +41 58 800 58 58

